

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C. 20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day month year) 12 October 2000 (12.10.00)	
International application No. PCT/NL00/00079	Applicant's or agent's file reference P48181PC00
International filing date (day month year) 09 February 2000 (09.02.00)	Priority date (day month year) 11 February 1999 (11.02.99)
Applicant GARSSEN, Gerrit, Jan et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
05 September 2000 (05.09.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer S. Mafla</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	--

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF THE RECORDING
OF A CHANGE(PCT Rule 92bis.1 and
Administrative Instructions, Section 422)

From the INTERNATIONAL BUREAU

To:

OTTEVANGERS, S., U.
Vereenigde
Nieuwe Parklaan 97
NL-2587 BN The Hague
PAYS-BAS

Date of mailing (day/month/year) 21 September 2001 (21.09.01)	IMPORTANT NOTIFICATION
Applicant's or agent's file reference P48181PC00	
International application No. PCT/NL00/00079	International filing date (day/month/year) 09 February 2000 (09.02.00)

1. The following indications appeared on record concerning:		
<input checked="" type="checkbox"/> the applicant	<input type="checkbox"/> the inventor	<input type="checkbox"/> the agent
<input type="checkbox"/> the common representative		
Name and Address STICHTING DIENST LANDBOUWKUNDIG ONDERZOEK Bornsesteeg 53 NL-6708 PD Wageningen Netherlands	State of Nationality NL	State of Residence NL
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	
2. The International Bureau hereby notifies the applicant that the following change has been recorded concerning:		
<input type="checkbox"/> the person	<input checked="" type="checkbox"/> the name	<input checked="" type="checkbox"/> the address
<input type="checkbox"/> the nationality		
<input type="checkbox"/> the residence		
Name and Address ID-LELYSTAD, INSTITUUT VOOR DIERHOUDERIJ EN DIERGEZONDHEID B.V. Edelhertweg 15 NL-8219 PH Lelystad Netherlands	State of Nationality NL	State of Residence NL
	Telephone No.	
	Facsimile No.	
	Teleprinter No.	
3. Further observations, if necessary:		
4. A copy of this notification has been sent to:		
<input checked="" type="checkbox"/> the receiving Office	<input type="checkbox"/> the designated Offices concerned	
<input type="checkbox"/> the International Searching Authority	<input checked="" type="checkbox"/> the elected Offices concerned	
<input type="checkbox"/> the International Preliminary Examining Authority	<input type="checkbox"/> other:	

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Dominique DELMAS Telephone No.: (41-22) 338.83.38
---	--

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference P48181PC00	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/NL 00/ 00079	International filing date (day/month/year) 09/02/2000	(Earliest) Priority Date (day/month/year) 11/02/1999
Applicant STICHTING DIENST LANDBOUWKUNDIG ONDERZOEK et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the language, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of the sequence listing:

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☒ furnished subsequently to this Authority in written form.

☒ furnished subsequently to this Authority in computer readable form.

☒ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☒ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

2. ☐ Certain claims were found unsearchable (See Box I).

3. ☐ Unity of invention is lacking (see Box II).

4. With regard to the title,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the abstract,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the drawings to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.



INTERNATIONAL SEARCH REPORT

Inter. Application No.

PCT/NL 00/00079

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 G01N33/68 C12Q1/37

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 G01N C12Q C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BELL, J. E. ET AL: "Prion protein immunocytochemistry - UK five center consensus report" NEUROPATHOL. APPL. NEUROBIOL. (1997), 23(1), 26-35 CODEN: NANEDL;ISSN: 0305-1846, XP002108042 the whole document	1-19
A	OESCH, BRUNO ET AL: "Properties of the Scrapie Prion Protein: Quantitative Analysis of Protease Resistance" BIOCHEMISTRY (1994), 33(19), 5926-31 CODEN: BICHAW;ISSN: 0006-2960, XP002108043 abstract	1,8
A	WO 98 32334 A (UNIV YALE ;MANUELIDIS LAURA (US)) 30 July 1998 (1998-07-30) abstract	1,8
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

11 May 2000

Date of mailing of the international search report

24/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Moreno, C



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/NL 00/00079

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 97 10505 A (UNIV CALIFORNIA) 20 March 1997 (1997-03-20) examples -----	1,8



INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 00/00079



Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9832334	A	30-07-1998	AU	5963598 A	18-08-1998
WO 9710505	A	20-03-1997	AU	707484 B	08-07-1999
			AU	7073596 A	01-04-1997
			CA	2231409 A	20-03-1997
			EP	0852011 A	08-07-1998
			JP	2000500005 T	11-01-2000
			NZ	318689 A	25-02-1999
			US	5846533 A	08-12-1998



PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference CJSO553	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US00/04904	International filing date (day/month/year) 25/02/2000	Priority date (day/month/year) 26/02/1999
International Patent Classification (IPC) or national classification and IPC C08F2/38		
Applicant THE UNIVERSITY OF AKRON et al.		
<p>1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of 8 sheets, including this cover sheet.</p> <p><input checked="" type="checkbox"/> This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</p> <p>These annexes consist of a total of 4 sheets.</p>		
<p>3. This report contains indications relating to the following items:</p> <p>I <input checked="" type="checkbox"/> Basis of the report</p> <p>II <input type="checkbox"/> Priority</p> <p>III <input type="checkbox"/> Non-establishment of opinion with regard to novelty, inventive step and industrial applicability</p> <p>IV <input checked="" type="checkbox"/> Lack of unity of invention</p> <p>V <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement</p> <p>VI <input type="checkbox"/> Certain documents cited</p> <p>VII <input type="checkbox"/> Certain defects in the international application</p> <p>VIII <input type="checkbox"/> Certain observations on the international application</p>		
Date of submission of the demand 21/09/2000	Date of completion of this report 29.06.2001	
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Barker, S Telephone No. +49 89 2399 8526 	

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/US00/04904

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17):*

Description, pages:

1-25 as originally filed

Claims, No.:

1-21 as received on 13/02/2001 with letter of 13/02/2001

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
☐ the language of publication of the international application (under Rule 48.3(b)).
☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
☐ filed together with the international application in computer readable form.
☐ furnished subsequently to this Authority in written form.
☐ furnished subsequently to this Authority in computer readable form.
☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/US00/04904

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

IV. Lack of unity of invention

1. In response to the invitation to restrict or pay additional fees the applicant has:

- ☐ restricted the claims.
- ☐ paid additional fees.
- ☐ paid additional fees under protest.
- ☐ neither restricted nor paid additional fees.

2. ☒ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.

3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is

- ☐ complied with.
- ☐ not complied with for the following reasons:

4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:

- ☐ all parts.
- ☐ the parts relating to claims Nos. .

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1-21
	No:	Claims	
Inventive step (IS)	Yes:	Claims	1-21
	No:	Claims	
Industrial applicability (IA)	Yes:	Claims	1-21
	No:	Claims	



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/US00/04904

2. Citations and explanations
see separate sheet



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US00/04904

Re. part IV

- 1). The amendments made by the applicant are considered to give rise to a lack of unity of invention for the following reasons. (Rule 68(1) PCT.) The subject matter of claim 1 (Invention I) is considered to define an invention which is distinct from that of claims 2-4 (Invention II). Claim 1 defines a polymerisation wherein the reducing agent is intended to regenerate the cobalt complex whereas in claims 2-4, no reducing agent need be present and the invention of claims 2-4 relates to the presence and structure of the halogen-containing polymers as defined in the polymerisation or method of forming functionalised polymer.

Re. part V

- 2). Novelty and inventive step is recognised for the subject matter of all of claims 1-21, for the reasons given below, so that the criteria set out in Art.33(2) and (3) are considered fulfilled.
 - a) US 5468785 A (D1) discloses a polymerisation process which comprises the step of photolyzing a cobalt(III) complex, in particular a cobaloxime complex, as defined in a partially polar solvent as defined. Monomer units are inserted at a free radical formed on the photolytic cleavage of a cobalt-carbon bond. Recombination of the growing polymer chain with the cobalt(II) complex, formed on photolysis, provides a pseudo-living polymerisation allowing, after a change of monomer, a site from which a linear block copolymer can be formed. It is reported that parallel to such mechanism, catalytic chain transfer occurs to provide a ethylenically unsaturated termination. Novelty is recognised for present claim 1 over D1 as there is no prior-art explicit disclosure of free-radical formation arising from reaction of the cobalt complex with a halogen-containing compound. Even if such reaction did occur between the cobalt(III) complex and the preferred chloroform solvent, then novelty is still recognised as D1 makes no mention of a reducing agent. The subject matter of independent claims 2,3 and 4 is also considered to be new as there appears to be no disclosure in D1 of a halogen-containing compound which is a polymer having a carbon-halogen bond as defined in any of these claims. Inventive step is further recognised as there appears to be no suggestion to make the relevant changes to arrive at the subject matter of the present claims.
 - b) US 5847060 A (D2) discloses the polymerisation of functionalised butadiene,



which can be a halo-substituted butadiene, using a free-radical initiator in the presence of a cobalt complex which acts as a chain transfer agent providing a substituted polybutadiene having terminal unsaturation. Novelty can be accepted for present claim 1 as there appears to be no disclosure of the polymerisation taking place in the presence of a reducing agent. Moreover, there appears to be no disclosure of the cobalt complex induced polymerisation of a halogen-functionalised polybutadiene as can be concluded from the information provided at example 8. That is, there does not appear to be a reaction wherein the cobalt complex reacts with a carbon-halogen bond to form a radical capable of participating in a catalytic chain transfer or pseudo-living polymerisation process. In particular it is noted that no fresh cobalt complex is added in example 8 of D2. As D2 does not suggest either the regeneration of cobalt complex by using a reducing agent or the cleavage by the reaction of a cobalt complex at a halogen-carbon bond on a halogen-substituted butadiene to provide a free-radical which can further polymerise, so the subject matter of present independent claims 1-4 and dependent claims 5-21 is considered not to be rendered obvious by the disclosure in D2.

c) US 5726263 A (D3) discloses the preparation of macromonomers by chain transfer catalysis (CTC) polymerisation methods which are then decolourised by selective extraction and/or adsorption. The CTC polymerisation of the (meth)acrylate monomer is carried out using an azo compound as initiator and a cobalt complex as chain transfer agent. In example 8, chloroform is present as a solvent. Present claim 1 is recognised to be new and not obvious as there appears to be neither disclosure or suggestion of the presence during polymerisation of a reducing agent. The subject matter of independent claims 2-4 is also recognised to be new and not obvious as there appears to be neither disclosure or suggestion of the CTC employing a polymer having halogen substitution. Indeed, as the problem in D3 is decolourising oligomers, the object of present claims 2-4 is well removed from that of D3. As all present independent claims are considered to be new and not obvious, so the subject matter of the dependent claims is also considered to be new and not obvious.

d) Chemical Abstracts, vol.132, abstract no.152175, relates to the published paper, SHIM, Anne K. et al: "The effect of solvent on the course of acrylate polymerizations mediated by cobaloxime and the development of novel initiating systems and novel multifunctional macromonomers" published in the journal



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US00/04904

Polymer Preprints, American Chemical Society, Division of Polymer Chemistry, 1999, 40(2), pp.132-133) (D4). The applicant has informed the IPEA that the D4 paper was published on 2nd August 1999 (02.08.1999) which is later than the date of filing of the US patent application (provisional application) 60/121895 filed on 26 February 1999 (26.02.1999). Inspection of the priority document by the IPEA concludes that present claim 1 is fairly based on the disclosure in the priority application so that D4 is not prior art in respect of that claim. The full claim to priority of claims 2-4 is less clearly established on account of the absence in the priority document of disclosure of polymers wherein the, "halogen atom is bound to an organic group that is pendent to the polymer backbone". Moreover disclosure in the priority document of "monomer" per se as used in claim 2 or of the acrylate monomer of claim 4 also appears to be lacking. Notwithstanding the seeming absence of a valid claim to priority, the subject matter of claims 2-4 is not rendered new or obvious by D4 as the only halogen compounds disclosed in D4 which react with cobaloxime compounds to form free radicals which allow polymerisation (or oligomerisation) of (meth)acrylate esters are halogenated solvents, in particular chloroform. Accordingly the lack of disclosure or suggestion in D4 of halogen-containing polymers leads to the conclusion that the subject matter of claims 2-4 and the dependent claims 5-21 meets the criteria of Arts.33(2) and (3) PCT.

e) Chemical Abstracts, vol. 128, Abstract no.154397, Arvanitopoulos, Labros D. et al, relates to "Photochemical polymerizations initiated and mediated by soluble organocobalt compounds", ACS Symp.Ser. (1998), 685 (Controlled Radical polymerization), 316-331 (D5). The disclosure concerns the polymerisation of acrylate esters by the use of organocobaloximes and a chloroform to provide a pseudo-living polymerisation. Present claim 1 is considered to be new and not obvious as there appears to be no mention of a reducing agent being used in the polymerisation. Present claims 2-4 are also considered to defined new and non-obvious subject matter as D5 appears to makes neither mention or suggestion of the halogen compound being a halogen-containing polymer as defined in claims 2,3 or 4. As all independent claims are considered to satisfy the criteria of Art.33(2) and (3) over D5, so the subject matter of dependent claims 5-21 also fulfil those criteria.



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US00/04904

Re. part VIII

- 3). a) The subject matter of claim 1 is considered vague, and therefore not clear, as it only defines the preparation of an admixture and gives no positive definition of a polymerisation step. Moreover, the claim is silent on the role of the reducing agent which forms the crux of the invention over the known prior art. (Art.6 PCT)
- b) The subject matter of claim 3 is unclear as it fails to repeat the recitation of the halogen being bound to an organic group pendent to the polymer background. Alternatively, it seems that any repetition of the position of the carbon-halogen bond in the polymers defined is unnecessary and would be better deleted. (Art.6 PCT)
- c) The wording of the description should be brought into accordance with the scope of the claims. (Art.6 PCT)

From the
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To:

PRINS, A.W.
VEREENIGDE

Nieuwe Parklaan 97
2587 BN The Hague

PAYS-BAS

1 APR 2001

TERMIJN

Beantwoord

Voorl.

def.

Bericht gezonden

aan

dd.

MAP

Applicant's or agent's file reference
P48181PC00

PCT

NRF 11-8-2001

NOTIFICATION OF TRANSMITTAL OF
THE INTERNATIONAL PRELIMINARY
EXAMINATION REPORT

(PCT Rule 71.1)

Date of mailing
(day/month/year)

05.04.2001

IMPORTANT NOTIFICATION

International application No.
PCT/NL00/00079

International filing date (day/month/year)
09/02/2000

Priority date (day/month/year)
11/02/1999

Applicant

STICHTING DIENST LANDBOUWKUNDIG ONDERZOEK et al.

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/



European Patent Office
D-80298 Munich
Tel. +49 89 2399 - 0 Tx: 523656 epmu d
Fax: +49 89 2399 - 4465

Authorized officer

Digiusto, M

Tel. +49 89 2399-8162





PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference P48181PC00	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/NL00/00079	International filing date (<i>day/month/year</i>) 09/02/2000	Priority date (<i>day/month/year</i>) 11/02/1999
International Patent Classification (IPC) or national classification and IPC G01N33/68		
Applicant STICHTING DIENST LANDBOUWKUNDIG ONDERZOEK et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 6 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 05/09/2000	Date of completion of this report 05.04.2001
Name and mailing address of the international preliminary examining authority: <div style="display: flex; align-items: center;"> <div> European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465 </div> </div>	Authorized officer Maucher, C Telephone No. +49 89 2399 7415





**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/NL00/00079

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-38 as originally filed

Claims, No.:

1-19 as originally filed

Drawings, sheets:

1/4-4/4 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
☐ the language of publication of the international application (under Rule 48.3(b)).
☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
☐ filed together with the international application in computer readable form.
☐ furnished subsequently to this Authority in written form.
☐ furnished subsequently to this Authority in computer readable form.
☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
☐ the claims, Nos.:



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/NL00/00079

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	2, 4-7, 9, 15-17
	No:	Claims	1, 3, 8, 10-14
Inventive step (IS)	Yes:	Claims	
	No:	Claims	2, 4-7, 9, 15-19
Industrial applicability (IA)	Yes:	Claims	1-19
	No:	Claims	

2. Citations and explanations
see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:
see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:
see separate sheet



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/NL00/00079

Point I:

The sequence listings filed on 26.4.2000 do not form part of the application as originally filed.

Point V:

Reference is made to the following documents:

- D1: NEUROPATHOL. APPL. NEUROBIOL.,
vol. 23, no. 1, 1997, p. 26-35
- D2: BIOCHEMISTRY,
vol. 33, no. 19, 1994, 5926-31
- D3: WO-A-97 10505

1. Article 33(2) PCT

- 1.1. The subject-matter of **independent claims 1 and 8 does not appear to be novel** (Article 33(2) PCT) in the light of the prior art, since the use of gdnSCN in tests reducing the risk of false positive results is known in the art.

D1, which is considered to be the closest prior art, already discloses the use of gdnSCN in hospitals (i.e. from patients; p 27, r col, 2 nd full paragraph, first lines) for treating a sample in a test for the presence or absence of insoluble prion protein (i.e. aberrant prion protein). Said protocol produced the clearest immunochemical results (i.e. the least false results; abstract).

The same applies to the following dependent claims:

- **claim 3** : D1, abstract: use of antibodies
- **claim 10** : D1, table 1: comparison of protocols comprising the use of gndSCN and of protocols omitting gndSCN
- **claims 11-13** : D1, table 2: antibody SP40 raised against a synthetic peptide based on prion disease protein gene sequence residues 219-232 (see specification, p 19, l 29 to p 20, l 6: residues 222-234 are in a



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/NL00/00079

- proteinase K resistant region of the sequence)
- **claim 14** : D1, table 2: SP40 (which is a functional equivalent of the claimed subject-matter)

- 1.2. The subject-matter of **dependent claims 2, 4** (see however VIII), **5-7, 9 and 15-17** is considered **novel** (Article 33(2) PCT), since none of the prior art documents discloses the claimed subject-matter.

However, the subject-matter of **dependent claims 2, 4** (see however VIII, 2.), **5** (D2, p 5927, r col 3rd full par; D3, p 8, l 10-14), **6-7** (D3, claim 2), **9, 15-16** (D2, p 5927, l col, 2nd par; D3, p 79, l 29 ff.), **17** (D3, claim 2) does not appear to be inventive (Article 33(3) PCT), since the additional features of said dependent claims are purely conventional and do not lead to an unexpected effect.

- 1.3. Although the subject-matter of **claims 18-19** is not defined by any technical feature (see VIII, 2.), a preliminary opinion can be given for novelty and inventive step:

If said claims were novel according to Article 33(2) PCT, they would not be inventive (Article 33(3) PCT) for the following reason. The mere packaging of substances known to be used in a known process does not involve any inventive activity and does therefore **not meet the requirements of Article 33(3) PCT**.

Point VII:

Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the documents D1-D3 is not mentioned in the description, nor are these documents identified therein.

Point VIII:

1. Claims 1, 8 and 14 in their present broad form are not supported by the description, since they are not commensurate with the contribution to the art of the



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/NL00/00079

present application (Article 6 PCT). Said objection is due to the inclusion of the unclear feature "functional equivalents thereof". No explicit example of a functional equivalent is given anywhere in the specification.

2. Claims 4 and 18-19 do not meet the requirements of Article 6 PCT in that the matter for which protection is sought is not clearly defined. The following functional statements do not enable the skilled person to determine which technical features are necessary to perform the stated function: "means for performing a method according to anyone of claims 8-17" (claim 18) and "for mass-screening purposes" (claims 4 and 19).





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : G01N 33/68, C12Q 1/37	A1	(11) International Publication Number: WO 00/48003 (43) International Publication Date: 17 August 2000 (17.08.00)
<p>(21) International Application Number: PCT/NL00/00079</p> <p>(22) International Filing Date: 9 February 2000 (09.02.00)</p> <p>(30) Priority Data: 99200391.3 11 February 1999 (11.02.99) EP</p> <p>(71) Applicant (for all designated States except US): STICHTING DIENST LANDBOUWKUNDIG ONDERZOEK [NL/NL]; Bornsesteeg 53, NL-6708 PD Wageningen (NL).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): GARSSSEN, Gerrit, Jan [NL/NL]; Prinses Marijkelaan 9, NL-3792 AJ Driebergen (NL). JACOBS, Jorg, Günther [NL/NL]; Horst 19-31, NL-8225 LV Lelystad (NL). LANGEVELD, Joannes, Pieter, Maria [NL/NL]; Stedenmeer 12, NL-3844 JB Harderwijk (NL). SMITS, Marinus, Adrianus [NL/NL]; Mastmeer 18, NL-3844-KE Harderwijk (NL). VAN KEULEN, Lucien, Johannes, Mattheus [NL/NL]; Kerkstraat 50, NL-8567 JH Oudemirdum (NL). SCHREUDER, Bram, Edward, Cornelis [NL/NL]; Rozengaard 13-10, NL-8212 DE Lelystad (NL). BOSSERS, Alexander [NL/NL]; Harderwijkstraat 137, NL-8244 DG Lelystad (NL).</p>	<p>(74) Agent: OTTEVANGERS, S., U.; Vereenigde, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>	
(54) Title: PRION TEST		
<p>(57) Abstract</p> <p>The invention is related to diagnostic methods for detecting transmissible spongiform encephalopathies (TSEs) such as BSE and scrapie and related disease in humans. The invention provides use of guanidine thiocyanate (gdnSCN) or a functional equivalent thereof for treating at least one sample derived from a mammal, including humans for reducing the risk of scoring a false-positive test result in testing said sample for the presence or absence of aberrant prion protein.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Title: Prion test

5 The invention is related to diagnostic methods for detecting transmissible spongiform encephalopathies (TSEs) such as BSE, scrapie and related diseases in animals and humans.

10 Bovine spongiform encephalopathy (BSE or mad cow disease) of cattle and scrapie of sheep are fatal, non-inflammatory neurodegenerative diseases caused by prions and are characterized by a long incubation period. In humans Creutzfeldt-Jakob disease (CJD), Gerstmann-Sträussler-Scheinker syndrome (GSS), fatal familial insomnia and kuru
15 belong to this category of TSEs.

Although scrapie, the prototype of the family of TSEs, in sheep and goats has been known for over 200 years (Pattison, 1988) and has been diagnosed world-wide (with the exception of New Zealand and Australia), it is only since
20 1986 that BSE has been described in cattle in the UK. By January 1998, there had been 170,259 confirmed cases of BSE in Great Britain and there may exist a great number of cases of not yet overt ("silent") BSE. BSE probably emerged because scrapie-contaminated sheep offal had been included in cattle
25 feeding-stuff via meat and bone meal and newly infected cattle material was then recycled (Wilesmith et al., 1991). This mechanism is quite plausible since ovine scrapie could be transmitted experimentally to several animal species, including cattle (Hourrigan, 1990; Gibbs, 1990).

30 Alternatively, recycling of offal from a rare case of spontaneous BSE for cattle feedstuff could also have led to the BSE epidemic. Moreover, the number of cattle in the UK with BSE reported annually is declining after the ban on feeding meat and bone meal in 1988.

Brain homogenates from cows with BSE produce after inoculation of mice a characteristic pattern of brain lesions in mice. Also characteristic incubation periods in inbred lines of mice are seen. This is identical to the pattern
5 elicited by brain tissue from individuals who recently have died from new-variant Creutzfeldt-Jakob disease (nvCJD; Bruce, 1997). The conclusion is that the BSE agent is identical to the nvCJD agent. Up to now, this variant has caused the death of 35 young Britons and one Frenchman (Will
10 et al, 1996; info: CJD Statistics per 30 November 1998, Internet).

There is also concern that the BSE strain that seems to be transmissible to humans may have infected sheep, where it could produce a disease hardly distinguishable from scrapie.
15 When its ominous strain-specific properties are maintained across the species barrier, sheep BSE may be a threat to human health, although scrapie by itself seems not to transmit to humans. Indeed, BSE agent has been transmitted experimentally to sheep by the oral route (Foster et al.,
20 1993) and thus could have the potential to infect sheep under field conditions. With the exception of a bioassay in mice, no diagnostic method is available to discriminate between BSE and scrapie in sheep at present.

Thus far, the only known component of the infectious
25 prion is an abnormal, disease-causing isoform of the "normal" prion protein (PrP) called PrP^{Sc} or aberrant prion protein. PrP, or normal prion protein, is ubiquitous in mammalian cells in a benign, cellular conformation (PrP^C) and is encoded within a single exon as a protein of about 250 amino
30 acid residues (figure 1). The PrP gene has been cloned and sequenced from a variety of species and there is a high degree of structural and organisational homology between mammalian PrP sequences (Schatzl et al., 1995). PrPs in many mammals have a 22-24 residues long N-terminal signal sequence

as well as a 22-24 residues long C-terminal signal sequence for attachment of a GPI-anchor. This glycosyl-phosphatidylinositol linkage is a fairly common means of anchoring proteins to membranes of eukaryotic cells. Further structural characteristics of the mature protein (of 206-210 amino acid residues) are one disulfide bond and two sites for Asn-linked glycosylation.

PrP^{Sc} originates from the normal cellular isoform (PrP^C) by a post-translational process since the amino acid sequence of PrP^{Sc} is identical to that predicted from cDNA or genomic nucleic acid sequences. Glycosylation patterns are also identical between PrP^C and PrP^{Sc}. Moreover, Caughey & Raymond (1991) demonstrated that PrP^{Sc} is made from a cell surface precursor that is identical to the normal PrP. PrP^{Sc} differs from the normal, membrane bound cellular prion protein by its relative protease resistance. Treatment with proteinase K (PK) for instance, results in complete proteolysis of PrP^C whereas in PrP^{Sc} the N-terminal part is removed before the amino acid at position 90 (human numeration). The protease-resistant core left is designated PrP27-30 after its electrophoretic behavior in SDS-PAGE as a protein molecule with $M_r = 27-30$ kDa, and this molecular species retains full infectivity.

Further distinguishing features of PrP^{Sc} are its thermal stability, a strong tendency to aggregate and insolubility in non-denaturing detergents, apparently connected with a different molecular structure. All attempts to identify a post-translational chemical modification that features in the conversion of PrP^C into PrP^{Sc} have been unsuccessful.

The lack of a molecular explanation for the observed differences between PrP^{Sc} and PrP^C led to the proposal that they must differ in conformation. Indeed, Fourier transform infrared spectroscopy detected a content of 43% of β -sheet and 30% of α -helix structure for purified hamster PrP^{Sc} and

an even higher β -sheet content of 54% for PrP²⁷⁻³⁰. On the other hand a low content of β -sheet structure and a high α -helix content of 42% was found in PrP^C, suggesting differences in secondary structure between the aberrant and
5 normal forms of PrP (Pan et al., 1993).

Due to its better solubility and the availability of recombinant forms of PrP^C, the three-dimensional structure of mouse PrP(121-231), involving three α -helices and a short antiparallel β -sheet, could be established by NMR (Riek
10 et al. 1996). In the mature murine PrP^C(23-231), this segment seems to have the same fold (Riek et al., 1997). Also the spatial structure of recombinant hamster PrP(29-231) has been examined (Donne et al., 1997).

A species barrier for prion infection has been
15 convincingly documented and found to vary widely depending on the pair of species involved and the direction of transmission. A structural basis for this species barrier is theoretically related to part or all of the amino acid replacements between the PrP of a given pair of species
20 (Billeter et al., 1997).

Within species, genetic polymorphism in the PrP gene has been found for example with mice, humans and sheep. In sheep amino acid substitutions in PrP at a few different positions were found to correlate with different predispositions for
25 the development of scrapie (Laplanche et al., 1993; Hunter et al., 1994; Belt et al., 1995; Bossers et al., 1996).

Studies of scrapie in goats and mice demonstrated reproducible variations in disease phenotype (length of incubation times and pattern of vacuolation) with the passage
30 of prions in genetically inbred hosts (Bruce & Fraser, 1991). The distinct varieties or isolates of prions were called "strains". Safar et al. (1998) made plausible that the biological properties of prion strains are enciphered in the conformation of PrP^{Sc} and that strains represent different

conformations of PrP^{Sc} molecules. Infection of Syrian hamsters with eight different hamster-adapted scrapie isolates produced PrP^{Sc} molecular species which, isolated from brains in the terminal stages of disease, differed with
5 respect to protease resistance and unfolding behavior under denaturing conditions. Differences in glycosylation have also been proposed as "strain-specific" properties (Collinge et al., 1996).

Animals and humans lack a TSE disease-specific immune
10 response and TSE diagnosis is based mainly on histopathological examination, which relies on the observation of neuronal degeneration, grey matter vacuolation (the spongiform change) and astrocytosis. A distinguishing feature of TSEs is the accumulation of aberrant protein
15 (PrP^{Sc}) in the brain under continuing biosynthesis of the normal cellular PrP^C. Species differences exist however, since the relative accumulation of PrP^{Sc} in brains of hamster and mouse is approx. 10x as high as in the ruminant. Unlike the normal PrP^C, PrP^{Sc} can aggregate into amyloid-like fibrils
20 and plaques and is a major component of brain fractions enriched for scrapie activity. Therefore, a more specific diagnosis of TSEs is detection of PrP^{Sc} either in situ e.g. by immunohistochemistry or in tissue homogenates e.g. by Western blot.

25 Several poly- or monoclonal antibodies to PrP have been described. The antisera were raised in mice, hamsters, rabbits and PrP null mice and as immunogens, peptides (as linear epitopes), purified and formic acid treated PrP^{Sc} from mice, hamster or sheep and recombinant PrP are being used.
30 However, except for one case (Korth et al., 1997), there have no antibodies been developed which can discriminate between native forms of PrP^C and PrP^{Sc}, and such antibodies cannot likely discern the difference between prion strains.

By Western blotting or immunohistochemistry PrP^{Sc} could be detected in sheep in brain, spleen, tonsil or lymph node material and even in a preclinical stage of scrapie (Schreuder et al., 1998). However, in BSE infected cattle PrP^{Sc} could not be detected outside the central nervous system, not even when clinical symptoms were present.

The intriguing mechanism of prion replication is not fully understood. According to the prevailing theory, the infectious PrP^{Sc} acts as a template in the replication of nascent PrP^{Sc} molecules. In other words PrP^{Sc} imposes its own conformation upon the cellular form PrP^C or an intermediate form. A thus far unknown protein X may function as a molecular chaperone in this formation of PrP^{Sc} (Prusiner et al., 1998).

Because of the connection between BSE and the nvCJD, and the possible transfer of BSE to other species including sheep, there is a need to monitor slaughter cattle and sheep for the presence of aberrant prion protein before the meat and meat products enter the human and animal food chain or into pharmaceuticals prepared for human and animal use. Mass screening of sheep and cattle should also be of help in view of eradication programmes of scrapie and BSE. Moreover, human blood and blood products may form a health threat on account of possible contamination with blood of CJD patients and the recent occurrence of the nvCJD. For these monitoring purposes a detection method for aberrant prion protein has to be developed which should be both fast, sensitive, reliable and simple.

Bioassays for PrP^{Sc} in which different doses of the analyte are administered to target animals, are generally regarded a gold standard but otherwise are cumbersome and costly. Moreover, their quantitative character is limited by a high variation. Immunohistochemical (IHC) approaches are

very useful insofar the presence of the analyte is directly made visible in the infected tissue. In particular testing said sample by histology or cytology allows a morphological comparison of healthy and diseased cells or tissue. Also the
5 presence of PrP^{Sc} can be indicated in a preclinical phase. However, and in general histological or cytological methods are not quantitative, and hardly applicable on a large scale.

For the diagnosis of TSEs founded on the demonstration of PrP^{Sc} in infected tissues and for the assessment of PrP^{Sc}
10 itself, several methods have been described and all are on an immunochemical basis. Most of these tests have been developed and used for research-like purposes, for instance in order to quantify PrP^{Sc} during purification procedures. In some cases calibration was with recombinant PrP (hamster or mice) or
15 with PrP^{Sc}, purified from scrapie-infected brains. Otherwise, responses were expressed as a function of mg tissue equivalents; in this way also sensitivity could be assessed by the minimum amount of tissue required for the PrP^{Sc} detection.

20 ELISA systems were designed for detection of PrP^{Sc}, isolated from brains of scrapie-affected mice and hamsters (Kascsak et al, 1987) and PrP^{Sc} from murine brain and spleen (Grathwohl et al., 1997). In these assays, the PrP^C fraction was beforehand removed by PK-treatment and the purified and
25 solubilised analyte was directly coated onto the microtiter plate. Solubilisation of PrP^{Sc} was by treatment with SDS or extraction with 77% formic acid, drying and resuspension in buffer (Kascsak et al. 1987). The denaturing action of formic acid was found to enhance the antibody response to PrP^{Sc}
30 considerably compared to untreated or SDS-treated material. In this ELISA rabbit antiserum to the mouse scrapie strain ME7 PrP^{Sc} was used.

Also successive solubilization of purified PrP^{Sc} by boiling in SDS, precipitation in cold methanol and sonication

in 3-4 M guanidine thiocyanate (gdnSCN) (Grathwohl et al., 1997) apparently enhanced coating-efficiency and/or epitope density under the denaturing action of gdnSCN. On the other hand, dissolving PrP^{Sc} in SDS appeared to inhibit adsorption of PrP^{Sc} onto the polystyrene microtiter plate. Although Grathwohl et al. (1997) state that their method could be a basis for a sensitive screening method for PrP^{Sc} in crude tissue extracts, their extraction and purification steps are impracticable and time-consuming (over 22 h). The sensitivity for brain tissue was such that PrP^{Sc} could be detected in 39 mg brain equivalents; the corresponding figure for spleen tissue amounted to 313 mg. Bell et al (1997), report comparative research of five research centres of in-house immunohistochemical methods for the detection of aberrant protein in CJD by histological staining of brain tissue sections. As to the use of gdnSCN, two of the five centres employ, in addition to formic acid, gdnSCN to pretreat their tissue sections to inactivate the prion agent to allow further processing of the tissues without the danger of infection. However, all over, the value of the addition of gdnSCN is questioned, and, in the opinion of one centre, it even increases background in histology. Effective decontamination of prion containing CJD material is also shown in WO98/32334.

A sandwich type of ELISA was used to monitor the bioproduction of recombinant hamster PrP(90-231), the protease resistant core of PrP^{Sc} (Mehlhorn et al., 1996). As a capture antibody the Fab fragment of mAb 3F4 was coated. This antibody was raised against hamster scrapie strain 263K and reacts with hamster, human and feline PrP. As the second antibody mAb 13A5 (to scrapie hamster PrP^{Sc}) was used. Samples from the different stages of purification were measured in this ELISA. However, the practical conditions under which PrP^{Sc}, in order to be detected as an antigen, is

brought into an unfolded state by chaotropic agents like 3-4 M gdnSCN, are not compatible with the immunochemistry of a sandwich type of ELISA.

Prusiner et al. (1990) used an enzyme-linked immunofiltration assay (ELIFA) which combines the properties of an immuno-dot blot and ELISA technique. By this method both PrP^C and PrP^{Sc} in scrapie brain homogenates of hamsters could be quantified against a standard curve of known amounts of purified hamster PrP27-30 (0.06-4 ng). Brain homogenates, diluted in buffer with 1 M gdnSCN and 0.05% Tween 20, were applied in 5 µl quantities to nitrocellulose membrane in a manifold filtration unit. Sequential steps for immunocomplex formation with mAb 13A5 and conjugation of enzyme were also done on this membrane. For detection, dots were cut out with a puncher and placed into a microtiter plate in which color was developed. Under these conditions, immunoreactivity of the dissociated and (partly) unfolded PrP^{Sc} is indistinguishable from that of PrP^C and in this way total PrP was measured. For the determination of the PrP^{Sc} fraction, the homogenate was treated with PK prior to the ELISA and PrP^C content was calculated by subtracting the PrP^{Sc} from the total PrP.

Oesch et al. (1994) refined this ELIFA method. Samples were applied on nitrocellulose filters in the ELIFA apparatus, procedures hereafter among which a 2h-preincubation in 4 M gdnSCN to render the aberrant protein sensitive to protease digestion, and substrate binding to mAb 13A5, up to and including binding with the enzyme were done on the membrane taken out of the apparatus. For detection, membranes were placed back in the ELIFA apparatus and reacted with substrate solution. Finally, the reaction mixture was pulled through into an ELISA plate placed underneath and colour development was measured. This whole procedure took over 20 hours.

Immuno-dot blotting was used by Serban et al. (1990) for the post mortem diagnosis of Creutzfeldt-Jakob disease in humans, scrapie in sheep and scrapie-infected hamsters and mice. Direct spotting of a rather impure analyte on e.g.

5 nitrocellulose filters instead of adsorption of a purified fraction of it onto the plastic surface of microtiter wells produces a more robust ELISA variant. This qualitative test was based on the intensified immunoreactivity of PrP^{Sc}-containing amyloid plaques after treatment with 3 M gdnSCN and the protease resistance of the PrP^{Sc} isoform.

10 Brains were extracted in detergent-containing lysis buffer and 4 µl amounts were spotted onto nitrocellulose membranes. Immunoreactivity of the spotted material after successive treatment with PK and 3 M gdnSCN was conclusive for the presence of PrP^{Sc} and confirmation of CJD and scrapie. Rabbit Ab R075 (to purified hamster PrP27-30) was able to detect PrP in the above four species. Out of a total of 28 human brain samples, 9 cases found positive by this method were also either defined as CJD or GSS by both

20 clinical diagnosis and a histopathological examination. For two cases, found positive by the blot procedure, histopathologic results were not available. The negative results of histopathology for CJD or GSS on the remaining 17 cases, coincided also with no indication for PrP^{Sc} with

25 the immuno-dot blot assay. In 12 histologically confirmed cases of natural scrapie in sheep, PrP^{Sc} was detected with the immunoblotting technique in the brains of 11 sheep. There are variations in the distribution of PrP^{Sc} in the brain of scrapie-affected sheep, since PrP^{Sc} was found in the spinal cord, cerebellum and pons/medulla of 2 sheep, but one sheep

30 also had PrP^{Sc} in the frontal and occipital cortex and the thalamus. This means that sampling of brain tissue could lead to false negatives due to regional variations in PrP^{Sc} content. The detection limit of this method for brain

extracts of scrapie-infected hamsters and mice ranged from 5-132 mg tissue equivalents, because these amounts still gave clearly visible spots. The duration of the test was, apart from an overnight incubation step, 6h.

5 Safar et al. (1998) developed a conformation-dependant fluorescent-ELISA that can discern various prion strains of hamsters. The assay detects a region of PrP^{Sc} that while exposed in normal PrP^C, becomes folded in the PrP^{Sc} molecule. Eu-labeled mAb 3F4 that reacts with a region of PrP^{Sc} only
10 after unfolding in 4 M gdnHCl and heating at 80°C for 5 min, was used in this assay. The immunoreactivity of the antibody to the denaturated region, as reflected by the fluorescence signal, is much higher than it is to PrP^{Sc} in its native form. The authors developed an algorithm which takes into
15 account that the immunoreactivity of antibody to denatured PrP in a sample of an affected brain is the summation of enhanced immunoreactivities of PrP^{Sc} and PrP^C during the transition from the native to the denatured states. Knowledge of the enhancement of immunoreactivity for PrP^C during
20 denaturation was a prerequisite for this approach. For this purpose calibration curves with different concentrations of purified PrP^C were constructed. It appeared that also PrP^C showed an enhanced immunoreactivity in 4 M gdnHCl compared to its native state, albeit in a moderate way ($\leq 1.8x$). From the
25 algorithm and the measurements of a native as well as a denatured sample, the content of PrP^{Sc} could be calculated. Although this method was validated for the determination of hamster brain, the authors aim at using it also for the detection of other mammalian prions, including human. In
30 order to improve the detection threshold of the assay they introduced an initial step to selectively precipitate PrP^{Sc} from raw material with sodium phosphotungstate. In combination with this sample pretreatment, the final sensitivity of the assay could be made high. The sensitivity

limit is less than or equal to 1 ng/ml (100 pg) of PrP^{Sc}. The test however, is still far from lending itself to large-scale use in view of too much labour and long incubation times.

Capillary electrophoresis was adapted by Schmerr et al. (1995, 1996, 1998a) as a diagnostic, immunochemical assay for scrapie. The authors claim a high sensitivity (approx. 135 pg PrP^{Sc}) of their test by measuring laser-induced fluorescence of a PrP-derived fluorescein-labeled peptide after its separation by free zone capillary electrophoresis. In a preceding competition step, this peptide was displaced from a preformed complex of the peptide and an antibody directed to the unlabeled peptide in competition with the analyte (PrP^{Sc}). Beforehand, PrP^C had been removed from the analyte solution by PK-treatment. The concentration of rabbit antiserum for complex-preformation was chosen so that the antibody would be limiting in the assay (adjustment to 50% of the maximum amount of immunocomplex). Four anti-(prion)-peptide antisera were prepared and evaluated. Assays using antisera to the peptides spanning mouse amino acid position 142-154 and 155-178, differentiated scrapie-positive sheep from normal animals. In spite of the high sensitivity of this method, sample processing is time-consuming (approx. 24h) and cumbersome since PrP^{Sc} from brain stem has to be concentrated and purified through steps like ultracentrifugation and HPLC.

Western blotting (WB) in combination with SDS-PAGE is also a suitable technique for diagnosis of TSEs and a variety of different extraction procedures and Western blotting methods has been described (Race et al., 1992; Beekes et al., 1995).

Usually, PrP^C is extracted from tissues with detergents that solubilize this membrane-bound protein in a mixed micelle. However, PrP^{Sc} in the presence of detergents, aggregates and therefore is not solubilized but can be spun down by ultracentrifugation. PrP^{Sc}-aggregates dissociate in monomers

under the denaturing conditions of heating in SDS solution with β -mercaptoethanol. In this way PrP^{Sc} is electrophoretically (SDS-PAGE) indistinguishable from PrP^{C} , unless a preceding treatment with PK has been applied. This
5 proteolytic treatment removes PrP^{C} and leaves $\text{PrP}27-30$, the truncated form of PrP^{Sc} .

Race et al. (1992) could find PrP^{Sc} in every brain of 8 sheep that were histologically positive for scrapie and even in brains of clinically positive sheep that were not
10 diagnosed as scrapie-positive by histology. For detection anti-peptide antibodies to residues 89-103 and 218-232 of the mouse PrP sequence were used. Apparently, the amount of tissue required to visualise PrP^{Sc} varied among sheep from <2 to 200 mg equivalents of brain tissue. Also PrP^{Sc} was
15 found in spleens and lymph nodes in 7 of 8 sheep that had the protease-resistant form detected in brain homogenates.

One method based on WB was officially approved by the European Union (EU) and the World Organisation for Animal Health (OIE) for BSE and scrapie diagnosis (Bradley et al.,
20 1994). A minimum amount of 2 mg equivalent of infected scrapie brain allows detection of the $\text{PrP}27-30$.

Above identified assays have never been used in large screening efforts for the detection of aberrant prion protein neither in animals nor in humans.

25

Thus far, two commercial assays have been announced. In 1997 the Swiss company Prionics Inc. launched its "BSE Western Test" intended for mass screening of slaughter cattle. A modified and optimised Western blot method was used
30 to detect the proteinase K-resistant $\text{PrP}27-30$ in bovine brain stem. For immunodetection mAb 6H4 was used, developed by immunizing PrP-null mice with recombinant bovine PrP. This antibody recognizes residues 147-155 of the bovine sequence as a linear epitope in native PrP^{C} and denatured PrP^{Sc} ; this

sequence is also recognised in sheep, human, pig and mouse. Incubation with anti-mouse IgG coupled to alkaline phosphatase and detection of the enzymatic product by chemiluminescence were the final steps of the assay. This
5 test requires an incubation step with PK and detects PrP²⁷⁻³⁰. Reliability is strengthened by the Western blot documentation of the decrease in size (internal control) of the prion protein from 30-33 to 27-30 kDa. The test can be done within hours and the expectation is that subclinical BSE
10 in post-mortem brains may be detected.

Also in 1997 the Irish Company Enfer Scientific Ltd. announced the development of a BSE post-mortem test. This immunoassay intended for mass screening uses a PrP antipeptide antiserum to detect PrP²⁷⁻³⁰ in samples of brain
15 tissue of cattle after removal of PrP^C by PK-treatment. Immunodetection was enhanced by chemiluminescence. Their claims are a result within 4 hours after receipt of samples and a capacity of 14,000 cattle a day and moreover, the catching of asymptomatic animals.

20 However, these two commercial tests, although claiming high sensitivity in detecting the aberrant protein, and thus claiming to have a low number of false-negative results, suffer from the low specificity associated with the claimed high sensitivity. When using the above tests one therefore
25 runs an increased risk of falsely identifying a negative sample as false-positive, thereby falsely identifying an animal as positive. For example, Switzerland slaughtered herds in which one or more cases of BSE had been confirmed. The "Swiss reference laboratory for animal TSE" examined the
30 brains of these 1761 apparently healthy cattle by an immunohistochemical method for signs of BSE and six positive cases were detected. Also Prionics Inc. tested these 1761 cattle brains by their "BSE Western Test". Four positive outcomes were identical to the ones found by the reference

laboratory, the other two were indicated as negative and moreover two other cattle were found positive by Western blotting. Thus a total of eight positive reactors were found, four of which overlapped. These eight were re-examined in the laboratory of Dr Kretzschmar (University of Göttingen) and in addition to the four undisputed cases, one of the two questionable cases identified by the reference laboratory could be confirmed (info: New Scientist, 1998, July 4 and Internet). Prionics for example scored 0.1% false-positives, indicating that in 1 of every thousand cases a sample causes a false-alarm due to false-positivity.

Tests scoring false-positive results (being in general not specific enough) have other consequences than tests scoring false-negative results (being in general not sensitive enough).

False-negative means that an in essence positive sample from a positive individual is scored negative, and thus is not suspected of having a TSE while in truth said individual is having a TSE. A false-negative diagnosis thus results in missing positive cases.

For humans, false-negative means that no diagnosis of TSE is made where said human actually has a TSE. This causes a wrong prognosis being established and wrong treatment being given, until a second test is done.

For animals, especially in those cases where slaughtered animals are tested, false-negative means that no diagnosis of TSE is made where said animal was actually infected and possibly capable of spreading the disease without having been noticed. Meat and other products from such a false-negative animal may contain aberrant prion protein. Such meat and meat products will be traded and eaten, and can thus be a source for further infection, notably of humans who even falsely trust that the animal has been tested well and the meat or meat product bears no risk.

False-positive means that an in essence negative sample from a negative individual is scored positive, and thus is at least suspected of having a TSE while in truth said individual is not having a TSE at all, but possibly another
5 condition.

For humans, false-positive means that a false diagnosis of TSE is made, here again resulting in false prognosis, and in faulty treatment. If said individual is not treated well as a consequence of the mis-diagnosis, his or her possible
10 other disease condition (the symptoms of which for example gave rise to the decision to test for TSE) receives no proper treatment.

For animals, false-positive means that a false diagnosis of TSE is made, however, since TSEs are notifiable diseases
15 that in general are met with strict eradication measures, said animal shall, at least in most Western countries be killed and destroyed. Furthermore, the herd from which said animals originated runs the same risk of being destroyed when the diagnosis is not corrected. For the slaughterhouse it
20 might mean that special laborious decontamination actions have to be implemented which mean temporary interference of use of the facilities and thus considerable loss of productivity. Additionally, the country where said animal or herd is falsely diagnosed for having a case of TSE among its
25 animals will be met with export restrictions. It goes without saying that, especially when said country has no (present) reported cases of TSE, such a false-positive diagnosis is highly detrimental for said countries position on foreign markets for animal products.

30 Understanding the above risks associated with false-negative or false-positive diagnoses becomes even more complicated when one understands that in general the level of false-positives scored by a diagnostic method or test is inversely related to the number of false-negatives scored by

the same test. It is an old diagnostic truth that, in many instances, a very sensitive test (having low numbers of false-negatives) cannot be very specific (and thus has a relative high number of false-positives) and vice versa.

- 5 However, and especially for mass screening tests that do not comprise histology or cytology, and wherein many samples need to be tested, tests having both high sensitivity and specificity are desired.

- 10 The invention provides use of guanidine thiocyanate (gdnSCN) or a functional equivalent thereof for treating at least one sample derived from a mammal for reducing the risk of scoring a false-positive test result in testing said sample for the presence or absence of aberrant prion protein, 15 in particular in testing said sample in a method other than histology or cytology. Using guanidine thiocyanate or its functional equivalents allows reduction of the signal arising from the normal prion protein (PrP^c) so that, for example when a gdnSCN-treated sample is compared with an untreated 20 sample, the PrP^c signal is greatly reduced. See for example Figure 3 herein describing the reduction of the signal from a PrP^{Sc} negative sample obtained by a use according to the invention. In a preferred embodiment, the invention provides use of guanidine thiocyanate (gdnSCN) or a functional 25 equivalent thereof for treating at least one sample derived from a mammal for reducing the risks of scoring both a false-positive test result or a false-negative test result in non-histologically or non-cytologically testing samples for the presence or absence of aberrant prion protein. In a preferred 30 embodiment, the invention provides use according to the invention in an immunoassay. The invention provides a reliable, simple and fast method, comprising use of gdnSCN or a functional equivalent thereof in a method for diagnosis of TSE being both highly specific as well as highly sensitive.

The invention also provides a method for reducing the risk of scoring a false-positive test result in testing a sample derived from a mammal for the presence or absence of aberrant prion protein comprising treating at least one
5 sample with gdnSCN or a functional equivalent thereof. Because of its simplicity and, due to its non-histological nature, speed, a method according to the invention particularly lends itself to mass screening purposes of e.g. post-mortem tissues in the slaughter-line of ruminants such
10 as cattle and sheep, but it is equally suitable in testing samples derived from other ruminants or experimental animals. In the human field the method could be used for e.g. screening lymphoid tissues and blood-derived products. Essentially, samples from all tissues, body fluids (e.g.
15 blood, liquor) and faeces can be used.

BSE or subclinical (silent) cases of BSE can for example be detected in samples automatically taken from the brain at the time that the heads are cut off from the slaughter-animals' trunk. The method can also be used in preclinical
20 stages during the development of scrapie, since tonsils which can be taken from the living animal, are proven to be an indicator tissue for preclinical scrapie and to contain PrP^{Sc} (Schreuder et al., 1998).

With scrapie in sheep as a model, we developed a method
25 to unambiguously distinguish PrP^{Sc} from PrP^C. This can be done on the basis of immunodetection of PrP^{Sc} without the need of a preceding elimination of PrP^C by enzymatic proteolysis (Figure 3).

In a modified embodiment, the invention provides use
30 according to the invention comprising use of a protease for treating said sample to reduce the presence of normal prion protein (Figure 4). This design is for example suited for the detection of aberrant prion protein in BSE.

Immunologically the signal of PrP^{Sc} can be enhanced in the presence of chaotropic agents. This enhancement is undoubtedly effected by dissociation of the polymeric PrP^{Sc} into oligo- or monomeric units, resulting in an increase of the number of epitopes available for the antibody. Moreover, the epitopes may be better exposed to the antibody through the protein defolding action of the denaturant.

Furthermore, the invention provides a method further comprising treating at least one first sample with gdnSCN or a functional equivalent thereof and leaving at least one second sample untreated with gdnSCN or a functional equivalent thereof and comparing the test result of said first sample with said second sample. We can for example discriminate between TSE-positive and negative cases after duplicate dot blotting of extracts of brain tissue onto a membrane. Extraction is in detergent-containing (lysis) buffer. One aliquot is left untreated and the other one is treated in 4 M gdnSCN (Figure 3) or, after PK-digestion, treated in 4 M gdnSCN (Figure 4). Then, after immunostaining, the treated sample is compared to the untreated sample. The signal of the normal protein (PrP^C) is greatly reduced or retains the same intensity upon treatment and can be further reduced by protease K digestion. In this way comparison of untreated and treated samples leads to a decrease or no increased of signal of samples from normal individuals but a significant increase of signal of samples from TSE-affected animals. By this dual internal control the discriminating value of the test is considerably enforced.

The invention also provides a method comprising immunological detection of said aberrant prion protein using at least one antibody directed against the aberrant protein, preferably directed against a proteinase K resistant part of the aberrant prion protein, for example wherein said antibody is directed against a proteinase K resistant N-terminal part

of the aberrant prion protein. Monoclonal or polyclonal antibodies can be used, possibly said antibody is raised against a peptide derived from the prion protein, for example wherein said peptide is selected from an N-terminal group consisting of residues 94-111 (like 94-105 and 100-111), a C-terminal group consisting of residues 222-234 and a group consisting of residues 145-177 (sheep numbering) or sequential homologues of the PK-resistant part of PrP^{Sc} (Figure 2) or functional equivalents thereof.

10 The invention also provides a method according to the invention wherein said protein is immunologically detected in an enzyme-linked immunoassay, for example wherein said enzyme-linked immunoassay comprises a dot blot assay.

15 Also, the invention provides a test kit having been provided with means for performing a method according to the invention. This kit for example contains a carrier matrix for spotting sample extracts from tissues, organs, cells or body fluids (e.g. nerve-tissue, blood cells, etc), buffers, solutions of gdnSCN and PK, primary antibody, enzyme-labeled
20 second antibody and enzyme-substrate. In a most preferred embodiment, said method or test kit is designed for mass-screening purposes.

25 The invention is further described in the detailed description herein without limiting the invention.

FURTHER DETAILED DESCRIPTION

MATERIALS AND METHODS

30 Phosphate buffered saline (PBS), pH 7.2 contained 136.89 mM NaCl, 2.68 mM KCl, 8.10 mM Na₂HPO₄ and 2.79 mM KH₂PO₄ in water.

PBTS: 0.2% (w/v) Tween-20 in PBS.

Two extraction buffers were used:

(a) 10 mM phosphate buffer, pH 7.0, 0.15 M NaCl and 0.25 M sucrose, used by Pan et al. (1992) to prepare microsomal fractions;

5 (b) lysis buffer (Collinge et al., 1996) consisted of 0.5% (w/v) Tergitol (type NP-40, nonylphenoxy polyethoxy ethanol, Sigma NP-40) and 0.5% (w/v) deoxycholic acid, Na-salt (Merck) in PBS, pH 7.2.

10 Guanidine thiocyanate (gdnSCN, purity > 99%; Sigma G 9277) solutions of 4 M were made up in water (pH 5.8).

Alkaline phosphatase-conjugated goat anti-rabbit IgG (GAR/AP) was from Southern Biotechnology Ass. (ITK,
15 Diagnostics B.V., Uithoorn).

Substrate for alkaline phosphatase was 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium (BCIP/NBT; tablets; Sigma B5655).

20

Usually, after PrP extraction, protease inhibitors were added to the extracts. (Complete, protease inhibitor cocktail tablets; Boehringer Nr. 1697498, Mannheim, Germany).

25 Proteinase K (EC 3.4.21.14, 20 units/mg lyophilisate Nr. 745723) and Pefabloc SC (4-(2-aminoethyl)-benzenesulfonyl fluoride, hydrochloride Nr. 1585916) were also from Boehringer. Incubation conditions for PrP-extracts with PK were 50 µg/ml enzyme for 30 min at 37°C. In order to stop
30 this enzymatic reaction, the incubation mixture was made 1 mM in Pefabloc added from a 100 mM stock solution of the inhibitor in water.

As a blocking agent nonfat dry milk (Protifar, Nutricia) was used.

A number of hydrophilic (14) and hydrophobic (5) membranes were tested as carrier matrix. A most successful
5 representatives, polysulphone or nitrocellulose membrane types were selected. Three membrane types were routinely used: nitrocellulose (NC) membrane with a 3 mm screen (Protran BA 85/21; 0.45 mm Nr. 405891) was from Schleicher & Schuell GmbH (Dassel, Germany), Immobilon-P,
10 (polyvinylidenedifluoride, PVDF) from Millipore B.V. (Etten Leur) and Zeta-Probe (quaternary amine-nylon membrane) was from BioRad.

An Ultra-Turrax T25 mixer with a 10 mm shaft (Ika Labortechnik GmbH, Staufen, Germany) was used to homogenize
15 brain tissue. The shaft was decontaminated in 1 M NaOH.

Water of 'Milli-Q' (Millipore) quality was used throughout.

20 *Primary antibodies*

These were intentionally designed for scrapie diagnosis. Antisera were induced in rabbits using synthetic peptides with sequences (12-mers) based on the sequence of ovine PrP protein. The sequences have such differences with the rabbit
25 PrP sequence that they induce not only antibodies which recognise these peptides but also the authentic PrP protein. Other animal species like mouse which have sequence differences could be as well suitable. The sequences used for immunisation were selected 12-mers from the protease K-
30 resistant domain of PrP^{Sc}. The selected 12-mer sequences represent domains that have a low tendency to form secondary structure (α -helix or β -sheet). The antisera are reactive in diagnostic dot blotting but also in Western blotting of both PrP^C and PrP^{Sc}, in ELISAs with as coated antigens the above

peptides or PrP protein, and in immunohistochemical detection. With the peptide derived from the ovine prion protein sequence 94-105, antisera R521 and R522 were produced in rabbits. Likewise sequence 100-111 yielded antisera R504, R505, R593, R594, R595, R596 and sequence 145-177 antiserum R532. The sequence 126-143 (ovine and bovine) gave rise to antiserum R568 while sequence 223-234 (ovine and bovine) yielded antisera R523 and R524. Peptides were synthesized and used to raise anti-peptide antisera in rabbits following previously published procedures (Van Keulen et al., 1995). Antisera were confirmed to be specific for sheep PrP (both undigested and after proteinase K treatment) on Western blots of partially purified prion protein from scrapie-affected sheep brain.

Sheep samples (brain stem, cervical spinal cord) were from scrapie-affected sheep, diagnosed by histopathological and immunohistochemical examination of the brain and from normal healthy sheep (Van Keulen et al., 1995). Samples from BSE-diagnosed cattle (histopathology, immunohistochemical examination and Western blotting) were from the cervical spinal cord or brain stem.

Procedure for immuno-dot blotting: 0.5 g portions of brain tissue were cut down with a scalpel and homogenized with an Ultra-Turrax mixer (20.000 rpm/15 sec) in 4.5 ml of ice-cold lysis buffer. The homogenates were centrifuged at 1000xg for 10 min or used without centrifugation as crude homogenate. If appropriate, an aliquot of the homogenate was incubated with PK at 37°C for 30 min after which the reaction was stopped with Pefabloc (1 mM). Otherwise a cocktail of protease inhibitors was immediately added to the homogenate. Suitable dilutions of the turbid supernatants or crude homogenates in lysis buffer were spotted in 1-3 µl amounts onto two blotting membranes and left for 15 min. One membrane was incubated in

4 M gdnSCN for 10 min, the other membrane was left untreated. Washing of the treated membranes was for 10 min in PBS on a rocking platform.

Membranes were blocked with 5% (w/v) Protifar in PBS for 5
1h at 20 °C and washed in PBTS with 1% (w/v) Protifar for 5
min at 20°C. A 1-2h incubation with the primary antibody
(1/1000 diluted in PBTS) at 20°C was followed by three
washing steps in PBTS for 5 min each. Next, the membranes
were incubated with AP-conjugated goat anti-rabbit IgG
10 (1/1000 diluted in PBTS) for 1-2 h at 20°C and washed in PBTS
three times for 5 min. Substrate solution was added and the
reaction was stopped with water.

RESULTS

15

DETECTION OF ABERRANT PRION PROTEIN IN SCRAPIE

Extraction efficiency for PrP^{Sc}

After homogenizing brain stem tissue of a scrapie-
20 affected sheep in extraction buffer (a) or in (b) (= lysis
buffer) and low-speed centrifugation which yielded
supernatant 1, aliquots of this supernatant were again
centrifuged at a higher speed (11,000xg, 10 min: 'high speed'
supernatant 1). The loose pellets left from the first
25 centrifugation step were adjusted with buffer to the original
volume, re-extracted and centrifuged at 1000xg, which yielded
a supernatant 2 and a loose pellet. In addition, aliquots of
all fractions were treated with PK.

1 µl extracts (diluted 1, 1/10 and 1/100x in their
30 respective buffers) were spotted onto NC and immunodetection
was with R522-7, an antiserum that has proven to detect ovine
PrP (Van Keulen et al, 1995).

For lysis buffer the highest signal intensity was
obtained for the supernatant 1. Compared to the results for

lysis buffer, the signals for extraction buffer (a) were lower for all fractions, except for the pellet. For fractions of the lysis buffer, decreased intensities were observed after pretreatment with proteinase K, especially for
5 supernatant 2, which indicates that this fraction is relatively enriched with PrP^C.

We observed dramatically intensified signals for the lysis buffer extracts, when these were diluted in 4 M gdnSCN. For supernatant 1, even after a 100-fold dilution, the signal
10 was clearly visible, which means that these scrapie brain stems PrP can be made visible in a tissue equivalent of 1 µg.

Divergent signal enhancement for PrP^{Sc} and PrP^C

Investigation of brain stem extracts of a scrapie-
15 negative sheep in lysis buffer revealed, even in a 80-fold dilution, clear signals of PrP^C. However, after pretreatment which PK no signal could anymore be observed. Surprisingly, instead of applying this PK-treatment, dilution of tissue extract in 4 M gdnSCN led also to a dramatic decrease of
20 signal intensity for PrP^C.

Next, instead of diluting lysis buffer extracted samples in 4 M gdnSCN, we applied serial dilutions of brain extracts of scrapie-positive and negative sheep in duplicate on NC membranes and incubated one membrane in 4 M gdnSCN for 10 min
25 while the other one was left untreated.

Immunodetection revealed that we easily could discriminate between scrapie positive (PrP^{Sc} and PrP^C) and scrapie negative (PrP^C) samples: a higher intensity with 4 M gdnSCN compared to an untreated sample means scrapie
30 positive, while a lower intensity with gdnSCN means scrapie negative.

This finding is the basis for a rapid and simple diagnostic test for TSEs. In this test there is in general no need for a preceding removal of PrP^C from the negative

sample.

Alternative denaturants and antisera

As an alternative for gdnSCN we investigated the effects
5 of other chaotropic agents. After dot blotting 3 μ l dilutions
of extracts of scrapie positive and negative brain stems,
separate NC membranes were incubated for 10 min in chaotropic
agents. The solutions used were: 4 M gdnSCN, 7.2 M urea, 4 M
KSCN, 1 M thiourea, NaOH (pH 11) in water and 98% formic
10 acid; besides one membrane was left untreated as a blank.
Results for immunodetection after KSCN and thiourea did not
differ from the blank. Urea induced a slight increase for the
scrapie positive material and formic acid enhanced the
intensity to the level of gdnSCN although this acid caused
15 considerable shrinking of the NC membrane. Optimum
enhancement with PVDF as a carrier was achieved by using 50%
formic acid, no membrane shrinkage was than observed. NaOH
(pH 11) on the other hand increased the signal for scrapie-
negative material.

20 Treatment with 4 M gdnSCN turned out to be the best
discrimination between scrapie positive and negative tissue
samples. Moreover, this effect appeared to be pH-invariant
since solutions of 4 M gdnSCN at pH 4 and 7 (in 50 mM
phosphate buffer), pH 6 (in water) and pH 9 (in 50 mM
25 carbonate buffer) gave identical results.

Five classes of antipeptide antisera to linear epitopes
of sheep PrP sequences (94-105, 100-111, 126-143, 145-177 and
223-234) were examined. For comparative reasons all sera were
used in a 1/500 dilution in PBTS. Antisera to the 94-105
30 sequence (R521, R522) and to the 100-111 sequence (R 505)
proved to have the best differentiating power. On the other
hand, with the antisera R568 and R532 to the sequences 126-
143 and 145-177 respectively, no immunoenhancing effect of
4 M gdnSCN on PrP^{Sc} could be detected.

Blotting membranes.

Comparison of results on NC membrane with those on Zeta-Probe showed for the latter a strong aspecific coloring of the entire membrane and consequently quaternary amine-nylon as a carrier was unsuitable. On the other hand, compared to nitrocellulose a stronger adsorption for PrP was shown for the PVDF membrane (Immobilon-P).

10 DETECTION OF ABERRANT PRION PROTEIN IN BSE

From brains of BSE-positive cattle, obtained from The Netherlands, the UK, Ireland, Belgium and Switzerland and of Dutch BSE-negative cattle (diagnosed by histopathology and immunohistochemical examination), brain stems were extracted with lysis buffer in the same manner as for sheep and the low-speed supernatant 1 was used for further examination. Brain stem extracts from confirmed scrapie-negative and positive sheep were used for comparison. Aliquots of extracts were also treated with proteinase K and 3 μ l amounts of dilutions in lysis buffer of PK-treated and untreated extracts were spotted onto NC membranes. Immunodetection was with 1/1000 dilutions of antisera to the 12-mer sequences 94-105 (antiserum R521), 100-111 (R505, R595, R596), 223-234 (R523, R524) and to the longer sequences 126-143 (R568) and 145-177 (R532).

Highest immunoreactivity was shown with antisera R505 and R595. After incubation with 4 M gdnSCN signal intensity of BSE-negative samples diminished; however, the immunoenhancing effect of 4 M gdnSCN on PrP^{Sc} in BSE-positive samples did not reach a comparable high level as for sheep PrP^{Sc} in scrapie. Surprisingly, antisera R523, R524 and especially R532 showed stronger immunoreactivity with bovine PrP^{Sc} than with PrP^C. Immunoreactivity of antisera R521 and R568 with bovine PrP

was very poor. No signal was obtained with the PK-treated material of BSE- and scrapie negative animals. PVDF showed a higher adsorption than NC membranes since immunostaining could be observed at higher dilutions on PVDF. The detection
5 limit of the test with sheep recombinant PrP spotted on PVDF and using antiserum R521 or R595 is about 50 pg. Using other detection methods, however, will of course result in even lower detection levels. Thusfar, 29 case of BSE and 131 negative controls were examined. The performance was 100%
10 (Table 1).

The design of one of our tests is that of a dot blot immunoassay which has an intrinsically higher sensitivity than an analogous ELISA assay in a microtiter plate, due to miniaturization within the blot and the higher binding
15 capacity of the matrix material (nitrocellulose, PVDF) than of a smooth polystyrene microtiter plate surface. Because of the divergent immunoreactivity of sheep PrP^C and PrP^{Sc} during denaturation, the discriminatory power for false positive samples of our test is much higher than that of the assay of
20 Safar and coworkers: in our assay the signal for PrP^C during denaturation in 4 M gdnSCN diminishes, whereas immunoenhancement (with 4 M gdnHCl) takes place in the assay of Safar and coworkers. As distinct from the test of Safar and coworkers, there is no need to calibrate our assay, it
25 can be performed within four hours and it lends itself to automation. Quantification will be with densitometric techniques. Other options for the design of our assay are an ELIFA-format combined with detection in solution of an enzyme-enhanced fluorescence or luminescence signal or time-
30 resolved detection of lanthanide fluorescence.

Table 1. Performance of dot-blot immunoassay on diagnosis of BSE

	true positive ¹	true negative ²	total	%	95% confidence interval ³
positive in test	29	0	29		
negative in test	0	131	131		
totals	29	131	160		
sensitivity				100	89.7 - 100
specificity				100	97.3 - 100

29

¹(Immuno)histochemically confirmed cases from five different countries. ²All confirmed negative. These 131 negative controls consisted of 45 cows suspected of BSE and 86 cows from a herd with a BSE-case. Likewise, the performance in the case of scrapie with samples of sheep is 100% (5 positive and 5 negative cases). ³ According to Blyth-Still-Casella.

REFERENCES

- Beekes, M., E. Baldauf, S. Caßens, H. Diringer, P. Keyes, A.C. Scott, A.H. Wells, P. Brown, C.J. Gibbs Jr & D.C. Gajdusek (1995). Western blot mapping of disease-specific amyloid in various animal species and humans with transmissible spongiform encephalopathies using a high-yield purification method. J. Gen. Virol. 76:2567-2576.
- Belt, P.B.F.M., I. Muileman, B.E.C. Schreuder, J. Bos-de Ruijter, A.L.J. Gielkens & M.A. Smits (1995). Identification of five allelic variants of the sheep PrP gene and their association with natural scrapie. J. Gen. Virol. 76:509-517.
- Bell J.E., Gentleman, S.M., Ironside, J.W., Mccardle, L., Lantos, P.L., Doey, L., Lowe, J., Ferguson, J., Luthert, P., McQuaid, S., and Allen, I.V. Prion protein immunocytochemistry - UK five centre consensus report. Neuropath. Appl. Neurobiol. (1997) 23:26-35.
- Billeter, M. R. Riek, G. Wider, S. Hornemann, R. Glockshuber & K. Wüthrich. (1997). Prion protein NMR structure and species barrier for prion diseases. Proc. Natl. Acad. Sci. USA 94:7281-7285.
- Bossers, A., B.E.C. Schreuder, I.H. Muileman, P.B.G.M. Belt & M.A. Smits (1996). PrP genotype contributes to determining survival times of sheep with natural scrapie. J. Gen. Virol. 77:2669-2673.
- Bradley et al. (1994). European Commission. Directorate General for Agriculture. Unit for Veterinary Legislation and Zootechnics. Protocols for the laboratory diagnosis and confirmation of bovine spongiform encephalopathy and scrapie. Report from the Scientific Veterinary Committee.
- Bruce, M.E., & H. Fraser (1991). Scrapie strain variation and implications. In: Chesebro B. (ed.). Transmissible spongiform encephalopathies. Current topics Microbiol.

Immunol. 172:125-138.

- Bruce, M.E., R.G. Will, J.W. Ironside, I. McConnell, D. Drummond, A. Suttle, L. McCardle, A. Chree, J. Hope, C. Birkett, S. Cousens, H. Fraser & C.J. Bostock (1997). Transmissions to mice indicate that 'new variant' CJD is caused by the BSE agent. *Nature* 389:498-501.
- Caughey, B. & G.J. Raymond (1991). The scrapie-associated form of PrP is formed from a cell surface precursor that is both protease- and phospholipase-sensitive. *J. Biol. Chem.* 266:18217-18223.
- Collinge, J., K.C.L. Sidle, J. Meads, J. Ironside & A.F. Hill (1996). Molecular analysis of prion strain variation and the aetiology of 'new variant' CJD. *Nature* 383:685-690.
- Donne, D.G., J.H. Viles, D. Groth, I. Mehlhorn, T.L. James, F.E. Cohen, S.B. Prusiner, P.E. Wright & H.J. Dyson (1997). Structure of the recombinant full-length hamster prion protein PrP(29-131): the N terminus is highly flexible. *Proc. Natl. Acad. Sci. U.S.A.* 94: 13452-13457.
- Foster, J., J. Hope & H. Fraser (1993). Transmission of bovine spongiform encephalopathy to sheep and goats. *Vet. Rec.* 133:339-341.
- Gibbs, C.J. Jr., J. Safar, M. Ceroni, A. DiMartino, W.W. Clark & J.L. Hourrigan (1990). Experimental transmission of scrapie to cattle. *Lancet* 335:1275.
- Grathwohl, K.-U. D., M. Horiuchi, N. Ishiguro & M. Shinagawa (1997). Sensitive enzyme-linked immunosorbent assay for detection of PrP^{Sc} in crude tissue extracts from scrapie-affected mice. *J. Virol. Meth.* 64: 205-216.
- Hourrigan, J.L. (1990). Experimentally induced bovine spongiform encephalopathy in cattle in Mission, Tex, and the control of scrapie. *J. Am. Vet. Med. Assoc.* 196:1678-1679.
- Hunter, N., W. Goldmann, G. Smith & J. Hope (1994). The association of a codon 136 PrP gene variant with the

- occurrence of natural scrapie. Arch. of Virol. 137:171-177.
- Ikegami, Y., M. Ito, H. Isomura, E. Momotani, K. Sasaki, Y. Muramatsu, N. Ishiguro & M. Shinagawa (1991). Pre-clinical and clinical diagnosis of scrapie by detection of PrP protein in tissues of sheep. Vet Rec. 128:271-275.
 - Kaszsak, R.J., R. Rubenstein, P.A. Merz, M. Tonna-DeMasi, R. Fersko, R.I. Carp, H.M. Wisniewski & H. Diringer (1987). Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins. J. Virol. 61:3688-3693.
 - Korth, C., B. Stierli, P. Streit, M. Moser, O. Schaller, R. Fischer, W. Schulz-Schaeffer, H. Kretzschmar, A. Raeber, U. Braun, F. Ehrensperger, S. Hornemann, R. Glockshuber, R. Riek, M. Billeter, K. Wüthrich & B. Oesch (1997). Prion PrP^{Sc}-specific epitope defined by a monoclonal antibody. Nature 390:74-77.
 - Laplanche, J.L., J. Chatelain, D. Westaway, S. Thomas, M. Dussausy, J. Brugere-Picoux & J.M. Launay (1993). PrP polymorphism associated with natural scrapie discovered by denaturing gradient gel electrophoresis. Genomics 15:30-37.
 - Mehlhorn, I., D. Groth, J. Stöckel, B. Moffat, D. Reilly, D. Yansura, W.S. Willett, M. Baldwin, R. Fletterick, F.E. Cohen, R. Vandlen, D. Henner & S.B. Prusiner (1996). High-level expression and characterization of a purified 142-residue polypeptide of the prion protein. Biochemistry 35:5528-5537.
 - Muramatsu, Y., A. Onodera, M. Horiuchi, N. Ishiguro & M. Shinagawa (1994). Detection of PrPres in sheep at the preclinical stage of scrapie and its significance for diagnosis of insidious infection. Arch. Virol. 134:427-432.
 - Oesch, B., M. Jensen, P. Nilsson & J. Fogh (1994). Properties of the scrapie prion protein: quantitative analysis of protease resistance. Biochemistry 33:5926-5931.
 - Pan, K.-M., N. Stahl, & S.B. Prusiner (1992) Purification and properties of the cellular prion protein from Syrian

hamster brain. Protein Science 1:1343-1352.

- Pan, K.-M., Baldwin, M., Nguyen, J., Gasset, M., Serban, A., Groth, D., Mehlhorn, I., Huang, Z., Fletterick, R.J., Cohen, F.E. & S.B. Prusiner (1993). Conversion of alpha-helices into beta-sheets features in the formation of the scrapie prion proteins. Proc. Natl. Acad. Sci. U.S.A. 90:10962-10966.
- Pattison, I.H. (1988). Fifty years with scrapie: a personal reminiscence. Vet. Rec. 123:661-666.
- Prusiner, S.B., M. Scott, D. Foster, K.-M. Pan, D. Groth, C. Mirenda, M. Torchia, S.-L. Yang, D. Serban, G.A. Carlson, P.C Hoppe, D. Westaway & S.J. DeArmond (1990). Transgenic studies implicate interactions between homologous PrP isoforms in scrapie prion replication. Cell 63:673-686.
- Prusiner, S.B., M.R. Scott, S.J. DeArmond & F.E. Cohen (1998). Prion protein biology. Cell 93:337-348.
- Race, R.E., D. Ernst, A. Jenny, W.Taylor, D.Sutton & B. Caughey (1992). Diagnostic implications of detection of proteinase K-resistant protein in spleen, lymph nodes, and brain of sheep. Am. J. Vet. Res. 53:883-889.
- Riek, R., S. Hornemann, G. Wider, M. Billeter, R. Glockshuber & K. Wüthrich (1996). NMR-structure of the mouse prion protein domain PrP(121-231). Nature 382:180-182.
- Riek, R., S. Hornemann, G. Wider, R. Glockshuber & K. Wüthrich (1997) NMR characterization of the full-length recombinant murine prion protein, mPrP(23-231). FEBS Lett. 413: 282-288.
- Safar, J., H. Wille, V. Itri, D. Groth, H. Serban, M.Torchia, F.E. Cohen & S.B. Prusiner (1998). Eight prion strains have PrP^{Sc} molecules with different conformations. Nature Medicine 10:1157-1165.
- Schatzl, H.M., M. Da Costa, L. Taylor, F.E. Cohen & S.B. Prusiner (1995). Prion protein gene variation among

- primates. J. Mol. Biol. 245:362-347.
- Schmerr, M.J., K.R. Goodwin, R.C. Cutlip & A.L. Jenny (1995). A competition assay to detect scrapie prion protein by capillary electrophoresis. J. Microcolumn Separations 7:521-527.
 - Schmerr, M.J., K.R. Goodwin, R.C. Cutlip & A.L. Jenny (1996). Improvements in a competition assay to detect scrapie prion protein by capillary electrophoresis. J. Chromatog. B 681:29-35.
 - Schmerr, M.J. & A. Jenny (1998). A diagnostic test for scrapie-infected sheep using a capillary electrophoresis immunoassay with fluorescence-labeled peptides. Electrophoresis 19:409-414.
 - Schreuder, B.E.C., L.J.M. van Keulen, M.E.W. Vromans, J.P.M. Langeveld & M.A. Smits (1996). Preclinical test for prion diseases. Nature 381:563.
 - Schreuder, B.E.C., L.J.M. van Keulen, M.E.W. Vromans, J.P.M. Langeveld & M.A. Smits (1998). Tonsillar biopsy and PrP^{Sc} detection in the preclinical diagnosis of scrapie. Vet. Rec. 142:564-568.
 - Serban, D., A. Taraboulos, S.J. DeArmond, S.J. & S.B. Prusiner (1990). Rapid detection of Creutzfeldt-Jakob disease and scrapie prion proteins. Neurology 40:110-117.
 - Van Keulen, L.J.M., B.E.C. Schreuder, R.H. Meloen, G. Mooij-Harkes, M. Poelen-van den Berg, M.E.W. Vromans & J.P.M. Langeveld (1995). Immunohistochemical detection and localization of prion protein in brain tissue of sheep with natural scrapie. Vet. Pathol. 32:299-308.
 - Van Keulen, L.J.M., B.E.C. Schreuder, R.H. Meloen, G. Mooij-Harkes, M.E.W. Vromans & J.P.M. Langeveld (1996). Immunohistochemical detection of prion protein in lymphoid tissues of sheep with natural scrapie. J. Clin. Microbiol. 34:1228-1231.

- Wilesmith, J.W., J.B.M. Ryan & M.J. Atkinson (1991). Bovine spongiform encephalopathy: epidemiological studies on the origin. Vet. Rec. 128:199-203.
- Will, R.G., J.W. Ironside, M. Zeidler, S.N. Cousens, K. Estibeiro, A. Alperovitch, S. Poser, M. Pocchiari, A. Hofman & P.G. Smith (1996). A new variant of Creutzfeldt-Jakob disease in the UK. Lancet 347:921-925.

LEGENDS TO THE FIGURES

Figure 1:

Amino acid sequences of human, rabbit, hamster, mouse, cattle and sheep PrP genes. The entire amino acid sequence of human PrP is given; open spaces in the other sequences indicate identity. Polymorphisms are indicated in bold at the top of each block and relate to the shaded positions. ↓: PHGGGWGQ. |: protease-sensitive site, right of which the sequence for the PK-resistant core of PrP^{Sc} is found. The mature PrP is devoid of N and C terminal signal peptides (in huPrP: amino acids 1-22 and 232-253, respectively).

human:

- Kretzschmar, H.A., S.B. Prusiner, L.E. Stowring & S.J. DeArmond (1986). Scrapie prion proteins are synthesized in neurons. *Am. J. Pathol.* 122:1-5.

rabbit:

- Loftus, B. & M. Rogers (1997). Characterization of a prion protein (PrP) gene from rabbit: a species with apparent resistance to infection by prions. *Gene* 184:215-219.
- Rubenstein, R., R.J. Kascsak, M. Papini, R. Kascsak, R.I. Carp, G. LaFauci, R. Melen & J. Langeveld (1998) *J. Neuroimmunology* (accepted).

golden Syrian hamster:

- Basler, K., B. Oesch, M. Scott, D. Westaway, M. Wälchli, D.F. Groth, M.P. McKinley, S.B. Prusiner & C. Weissmann (1986). Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene. *Cell* 46:417-428.

mouse:

- Locht, C., B. Chesebro, R. Race & J.M. Keith (1986). Molecular cloning and complete sequence of prion protein cDNA from mouse brain infected with the scrapie agent. Proc. Natl. Acad. Sci. USA 83:6372-6376.
- Westaway, D., P.A. Goodman, C.A. Mirenda, M.P. McKinly, G.A. Carlson & S.B. Prusiner (1987). Distinct prion proteins in short and long scrapie incubation period mice. Cell 51:651-662.

cattle:

- Goldmann, W., N. Hunter, T. Martin, M. Dawson & J. Hope (1991). Different forms of the bovine PrP gene have five or six copies of a short, g-c-rich element within the protein-coding exon. J. Gen. Virol. 72:201-204.

sheep:

- Goldmann, W., N. Hunter, J.D. Foster, J.M. Salbaum, K. Beyreuther & J. Hope (1990). Two alleles of a neural protein gene linked to scrapie in sheep. Proc. Natl. Acad. Sci. USA 87:2476-2480.

Figure 2:

Peptide sequences, derived from the prion protein structures of six species (**hu**=human, **rb**=rabbit, **ha**=hamster, **mo**=mouse, **bo**=cattle, **ov**=sheep). The amino acid sequence of the human peptides is given; open spaces in the other sequences indicate identity.

Antipeptide antibodies were raised in rabbits against the peptides of the ovine structure. Corresponding antisera are indicated **R5xx** at the top of each set of sequences.

Figure 3:

Prion test in which the extract was applied to two pieces of NC-membrane indicated: untreated and treated. A 10% (wt/vol.%) extract of brain-stem tissue was 1/3 diluted and 1 μ l applied to NC-membrane. Then, one piece of membrane was incubated in solution without gdnSCN (untreated), the other was incubated in 4 M gdnSCN-containing solution (treated). Further incubations for immunochemical visualization with first antibody (R522-7) and alkaline-phosphatase conjugate were according to standard procedures. In each 3-fold dilution series, the first (left) spot represents 33 μ g of tissue equivalents.

Figure 4:

Prion test, in which the extract was applied to two pieces of PVDF-membrane indicated: untreated and treated. For each negative and positive case, a 10% (wt/vol.%) extract of brain-stem tissue was prepared and divided in two portions, of which one was incubated with proteinase K (PK-digested extract) or not (undigested extract). Next, each of the extracts was 1/3 diluted and 3 μ l applied to PVDF-membrane. Then, the piece of membrane with the undigested extract was incubated in solution without gdnSCN (untreated), the membrane with digested extract was incubated in 4 M gdnSCN-containing solution (treated). Further incubations for immunochemical visualization with first antibody (R595-4) and alkaline-phosphatase conjugate were according to standard procedures. In each 3-fold dilution series, the first (left) spot represents 100 μ g of tissue equivalents.

Claims

1. Use of guanidine thiocyanate (gdnSCN) or a functional equivalent thereof for treating at least one sample derived from a mammal for reducing the risk of scoring a false-positive test result in testing said sample for the presence or absence of aberrant prion protein.
2. Use according to claim 1 for reducing the risk of scoring a false-negative test result.
3. Use according to claim 1 or 2 wherein said sample is tested in a immunoassay.
4. Use according to anyone of claims 1 to 3 wherein said immunoassay is designed for mass-screening purposes.
5. Use according to anyone of claims 1 to 4 further comprising use of a protease for treating said sample to reduce the presence of normal prion protein.
6. Use according to anyone of claims 1 to 5 wherein said sample is derived from a ruminant.
7. Use according to claim 6 wherein said ruminant is ovine or bovine.
8. A method for reducing the risk of scoring a false-positive test result in testing a sample derived from a mammal for the presence or absence of aberrant prion protein comprising treating at least one sample with gdnSCN or a functional equivalent thereof.
9. A method according to claim 8 further comprising reducing the risk of scoring a false-negative result.
10. A method according to claim 8 or 9 further comprising treating at least one first sample with gdnSCN or a functional equivalent thereof and leaving at least one second sample untreated with gdnSCN or a functional equivalent thereof and comparing the test result of said first sample with said second sample.
11. A method according to anyone of claims 8 to 10 further comprising immunological detection of said aberrant prion

protein using at least one antibody directed against a proteinase K resistant part of the aberrant prion protein.

12. A method according to claim 11 wherein said antibody is directed against a proteinase K resistant N-terminal part of
5 the aberrant prion protein.

13. A method according to claim 11 or 12 wherein said antibody is raised against a peptide derived from the prion protein.

14. A method according to claim 13 wherein said peptide is
10 selected from the groups listed in figure 2 or functional equivalents thereof.

15. A method according to anyone of claims 11 to 14 wherein said protein is immunologically detected in an enzyme-linked immunoassay.

15 16. A method according to claim 15 wherein said enzyme-linked immunoassay comprises a dot-blot assay.

17. A method according to any one of claims 8 to 16 wherein said mammal is ruminant, preferably wherein said ruminant is ovine or bovine.

20 18. A test kit having been provided with means for performing a method according to anyone of claims 8 to 17.

19. A test kit according to claim 18 which is designed for mass-screening purposes

1	hu	M A - - N L G C W M L V L F V A T W S D L G L C K K R P K P -	28
1	rb	- - - H Y Y L L A A M T V V - G	29
1	ha	- - - S Y Y L A T M T V V - -	28
1	mo	- - - G Y Y L A T M T V V - -	28
1	bo	V K S H I S I M M V V G	31
1	ov	V K S H I S I M V G	31
29	hu	G G W N T G G S R Y P G Q G S P G G N R Y P P Q G G G G G W G Q P	60
30	rb	- - - - - S - - - - -	60
29	ha	- - - - - - - - - - -	60
29	mo	- - - - - - - - - - -	59
32	bo	- - - - - - - - - - -	63
32	ov	- - - - - - - - - - -	63
61	hu	H G G G W G Q P H G G W G Q P H G G G W G Q P H G G G - W G	90
61	rb	- - - - - - - - - - -	90
61	ha	- - - - - - - - - - -	90
60	mo	- - - - - S - - - - -	89
64	bo	- - - - - - - - - - -	94
64	ov	- - - - - G - - - - -	94
91	hu	Q G G G G T H S Q W N K P S K L P K T N F K H M A G A A A A G A	120
91	rb	- - - - - N N G - - - - -	119
91	ha	- - - - - N N - - - - -	120
90	mo	- - - - - N N - - - - -	119
95	bo	- - - - - G - - - - -	123
95	ov	- S - - - - - V - - - - -	123
121	hu	V V G G L G G Y M L G S A M S R P F I H F G S D Y E D R Y Y	150
120	rb	- - - - - - - - - - -	149
121	ha	- - - - - - - - - - -	150
120	mo	- - - - - M - - - - -	149
124	bo	- - - - - I - - - - -	153
124	ov	- - - - - L - - - - -	153
151	hu	H R E N M H R Y P N Q V Y Y R P M D Q E Y S N Q N N F V H N C V	180
150	rb	- - - - - Y - - - - -	179
151	ha	- - - - - N Y - - - - -	180
150	mo	- - - - - Y - - - - -	179
154	bo	- - - - - V - - - - -	183
154	ov	- - - - - Y - - - - -	183
181	hu	N I T I K Q H T V V T T T T K G E N S F T E T D V K M M E R V V	210
180	rb	- - - - - V - - - - -	209
181	ha	- - - - - - - - - - -	210
180	mo	- - - - - T - - - - -	209
184	bo	- - - - - V E - - - - -	213
184	ov	- - - - - V - - - - -	213
211	hu	E Q M C I T Q Y E R E S Q A Y Y - Q R G - S S M - V - L F S S P P V	240
210	rb	- - - - - Q Q - - - - -	239
211	ha	- - - - - T Q Q - - - - -	241
210	mo	- - - - - V Q Q - - - - -	241
214	bo	- - - - - Q - - - - -	243
214	ov	- - - - - Q - - - - -	243
241	hu	I L L I S F L I F L I V G	253
240	rb	- - - - - - - - - - -	252
242	ha	- - - - - M - - - - -	254
242	mo	- - - - - - - - - - -	254
244	bo	- - - - - - - - - - -	256
244	ov	- - - - - - - - - - -	256

Amino acid sequences of human, rabbit, hamster, mouse, cattle and sheep PrP genes. The entire amino acid sequence of human PrP is given; open spaces in the other sequences indicate identity. Polymorphisms are indicated in bold at the top of each block and relate to the shaded positions. ↓: PHGGGWGQ. | : protease-sensitive site, right of which the sequence for the PK-resistant core of PrP^{Sc} is found. The mature PrP is devoid of N and C terminal signal peptides (in huPrP: amino acids 1-22 and 232-253, respectively).

Fig. 1

313 Rec'd PCT/PTO 10 AUG 2001

Rabbit anti-peptide antisera to parts of the ovine PrP-structure

R521, R522 (to ovine sequence 94-105)

90	hu	G	Q	G	G	T	H	S	Q	W	N	K	P
90	rb	G			-			N		G			
90	ha	G						N					
89	mo	G						N					
94	bo	G			-			G					
94	ov	G			-	S							

R504, R505, R593-596 (100-111)

97	hu	S	Q	W	N	K	P	S	K	P	K	T	N
96	rb	N			G							S	
97	ha	N											
96	mo	N											
100	bo	G											
100	ov												

R568 (126-143)

123	hu	G	G	L	G	G	Y	M	L	G	S	A	M	S	R	P	I	I	H
122	rb																L		
123	ha																M	M	
122	mo												M				M		
126	bo												L				L		
126	ov												L				L		

R532 (145-177)

142	hu	G	S	D	Y	E	D	R	Y	Y	R	E	N	M	H	R	Y	P	N	Q	V	Y	Y	R	P	M	D	E	Y	S	N	Q	N	N
141	rb		N												Y											V		Q						
142	ha		N		W										N											V		Q			N			
141	mo		N		W										Y											V		Q						
145	bo																									V		Q						
145	ov		N												Y											V		R						

R523, R524 (223-234)

220	hu	R	E	S	Q	A	Y	Y	-	Q	R	G	-	S
219	rb	Q				A			-			-	A	-
220	ha	K						D	G		R	-	A	-
219	mo	K						D	G		R	-	A	-
223	bo							-					A	
223	ov							-					A	


Fig. 2

518 Rec'd PCT/PTO AUG 2001

Fig. 3

example of test-result with brain-stem extract
from sheep

SHEEP







510 Rec'd POT/PT 0 AUG 2001





Fig. 4

example of test-result with brain-stem tissue
from sheep and cattle

CATTLE

	untreated	treated (PK, gdnSCN)
BSE negative		
BSE positive		
	3-fold sample dilution	3-fold sample dilution

SHEEP

	untreated	treated
scrapie negative		
scrapie positive		
	3-fold sample dilution	3-fold sample dilution

518 Rec'd PCT/ 10 AUG 2001